

Glypure™

Cosmetic-Grade Glycolic Acid

Formulation—Cuticle Cream

	INCI Name	Trade Name	Supplier	wt%
Phase A	Purified Water USP	Purified Water		40.00
	Magnesium Aluminum Silicate	Magnabrite S	Amcol	0.60
	Glycerin 99% USP	Glycerin USP	Spectrum	3.00
	Propylene Glycol	Propylene Glycol USP	Spectrum	1.50
	Xanthan Gum NF	Keltrol CG-T, Vanzan NF	Kelco, R.T. Vanderbilt	0.30
	Disodium EDTA	Versene NA2X, Dissolvine NA2S	Dow, Akzo	0.05
	Phase B	Purified Water	Purified Water	
	Glycolic Acid (70%)¹	Glypure™ (70%)	Chemours	5.75
	Triethanolamine 99% NF ²	Trolamine 99	Dow, Ineos	0.90
	To pH 3.3–3.5 ³			
Phase C	Glyceryl Stearate/PEG-100 Stearate	Arlacel 165V	Croda	3.50
	Glyceryl Stearate	Cutina GMS V	BASF	0.50
	Stearic Acid	Emersol 7036	Emery Oleochemicals	2.00
	Cetearyl Alcohol	Lanette O	BASF	2.50
	White Petrolatum	Ultima USP	Penreco	4.00
	Tocopheryl Acetate	Vitamin E Acetate	BASF	0.25
	Octyldodecyl Myristate	ODM 100	Barnet	2.50
	Octyldodecyl Neopentanoate	Elefac I-205	Alzo	3.00
	Dimethicone	DC 200/100 cst	Dow Corning	1.00
	Ethylhexyl Hydroxystearate	Wickenol 171	Alzo	3.00
	Phase D	Cyclomethicone	ST-Cyclomethicone 5-NF	Dow Corning
	Isododecane	Permethyl 99A	Presperse	1.50
	Isohexadecane	Permethyl 101A	Presperse	1.50
Phase E	Butylene Glycol (and) Glycerin (and) Chlorphenesin (and) Methylparaben	Elestab FL-15	Lab.Serobiologiques/ BASF	2.50
Phase F	Dye, Fragrance, and Additives ⁴	As Desired	As Desired	
	Purified Water	Purified Water		qs to 100%

Notes:

¹Glypure™ (99%) may be substituted for Glypure™ (70%). Compensate the Purified Water percentage accordingly.

²May use other suitable alkalis, e.g., Potassium Hydroxide, Ammonium Hydroxide, or Sodium Hydroxide.

³Do not exceed 2.5% of Triethanolamine to comply with EU regulations. If necessary, add another neutralizing agent.

⁴Compensate the Purified Water percentage accordingly for any additives.

Manufacturing Procedure

- 1) Prepare Phase A by adding water to the main vessel and begin mixing.
- 2) Slowly add A2 and mix for 15–30 min to hydrate.
- 3) Pre-wet A5 in A3 and A4, and mix until a uniform dispersion is obtained. Add to A1 and A2, and mix until uniform.
- 4) Add A6, and continue mixing while heating to 70–75 °C (158–167 °F).
- 5) In a separate vessel, mix B1 and B2 (Glypure™) and adjust pH accordingly with B3.
- 6) In a separate vessel, add C1 through C10, and heat to 70–75 °C (158–167 °F). Begin mixing slowly when solid ingredients begin to melt.
- 7) When Phases A and C are at the proper temperature range, add Phase C to Phase A slowly. When complete, homogenize for 5 min.
- 8) Begin cooling. When Phase AC is at 62–65 °C (144–149 °F), pre-blend D1 through D3 and add to Phase AC. Homogenize while cooling is continued.
- 9) When Phase ACD is 45–50 °C (113–122 °F), add pH-adjusted Phase B. Continue homogenizing and cooling.
- 10) Continue cooling and add E and F at 36–38 °C (97–100 °F) and any make-up water due to additive additions and water losses.
- 11) Adjust to pH 3.7–4.0 if necessary with appropriate neutralizing agent.
- 12) Turn off homogenizer and sweep mix to <30 °C (86 °F).

For more information, visit glycolicacid.chemours.com or call (800) 441-9593.

The information set forth herein is furnished free of charge and based on technical data that Chemours believes to be reliable. It is intended for use by persons having technical skill, at their own risk. Because conditions of use are outside our control, Chemours makes no warranties, express or implied, and assumes no liability in connection with any use of this information. Nothing herein is to be taken as a license to operate under, or a recommendation to infringe, any patents or patent applications.

© 2020 The Chemours Company FC, LLC. Glypure™ is a trademark of The Chemours Company FC, LLC. Chemours™ and the Chemours Logo are trademarks of The Chemours Company.

Replaces: K-17605

C-11797 (3/20)