

Glypure™

Cosmetic-Grade Glycolic Acid

Formulation—Skin Care Day Cream with Sunscreen



Glypure™ penetrates the skin efficiently—readjusting water percentages in the epidermis, stimulating collagen synthesis, and promoting cell turnover. It is also an efficient pH adjuster.

- Improves the look and feel of skin
- Promotes exfoliation
- Improves skin texture
- Reduces the appearance of fine lines and wrinkles
- Improves the appearance of sun-damaged skin
- Fights the signs of aging
- Helps even out skin tone

Phase	Trade Name	Wt%	INCI Name	Supplier
A1	Purified Water USP	30.00	Purified Water USP	
A2	Veegum® Ultra	1.00	Magnesium Aluminum Silicate	Vanderbilt Minerals, LLC
A3	Glycerin USP	3.00	Glycerin 99% USP	Spectrum Chemical
A4	Keltrol® CG-T	0.30	Xanthan Gum	CP Kelco
A5	Edeta® BD	0.20	Disodium EDTA	BASF
A6	Triethanolamine 99%	1.50	Triethanolamine 99% NF	–
B1	Purified Water USP	8.00	Purified Water USP	
B2	Glypure™	11.50	Glycolic Acid (70%)¹	Chemours
B3	Ammonium Hydroxide Solution	1.5-2.0	Ammonium Hydroxide 28%–to pH 3.8 or desired pH	–
C1	Emersol® 7036	3.75	Stearic Acid	Emery Oleochemicals
C2	Alfol 16	1.50	Cetyl Alcohol	Sasol
C3	Arlacel™ 165	4.00	Glyceryl Stearate/PEG-100 Stearate	Oxiteno
C4	Escalol® 557	7.50	Octylmethoxycinnamate	Ashland
C5	Captex® 200	1.00	Propylene Glycol Dicaprylate/Dicaprate	Abitec
C6	Vitamin E Acetate, USP	0.50	Tocopherol Acetate	Spectrum Chemical
C7	Eutanol® G-16	4.00	Isocetyl Alcohol	BASF
C8	Xiameter® PMX-200 Silicone Fluid 100CS	2.00	Dimethicone	Dow Corning
C9	Crodamol™ AB	3.00	C12-15 Alkyl Benzoate	Croda
C10	Elefac™ I-205	5.00	TiO ₂ /Isononyl Isononanoate/Stearic Acid/Aluminum Hydroxide	Alzo International
C11	Tioveil™ 50 FCM	5.00	Titanium Dioxide (and) Cyclomethicone (and) C12-15 Alkyl Benzoate (and) Polyhydroxystearic Acid (and) Aluminum Stearate (and) Alumina	Croda
D1	Phenonip®	1.00	Phenoxyethanol (and) Methylparaben (and) Ethylparaben (and) Butylparaben (and) Propylparaben (and) Isobutylparaben	Clariant
E1	As Desired	0.00	Fragrance ²	As Desired
qs	Purified Water USP	qs to 100%	Purified Water USP	

Notes:

¹Glypure™ (99%) may be substituted for Glypure™ (70%). Compensate the purified water percentage accordingly.

²Compensate the purified water percentage accordingly for any additives.

In lieu of Glypure™, formulators and manufacturers must use Glypure™ L for products used or distributed in Canada or Australia and in Europe for nail care products.

Manufacturing Procedure

1. Prepare phases A, B, and C separately.
2. When preparing phase C, homogenize for 30 min after heating and addition of the TiO₂ dispersion.
3. Heat phases A and C to 70-75 °C (158-167 °F).
4. Blend phases A and C, stirring and homogenizing for a minimum of 10 min.
5. Cool to 65 °C (149 °F).
6. Add phase B and stir with homogenizing for a minimum of 10 min.
7. Cool and add preservative, fragrance, and purified water below 40 °C (104 °F).

Glypure™ has proven benefits in hair, skin, and nail care formulations. To learn more about the benefits of Glypure™, visit www.glypure.com.

For more information, visit glycolicacid.chemours.com or call (800) 441-9593.

The information set forth herein is furnished free of charge and based on technical data that Chemours believes to be reliable. It is intended for use by persons having technical skill, at their own risk. Because conditions of use are outside our control, Chemours makes no warranties, expressed or implied, and assumes no liability in connection with any use of this information. Nothing herein is to be taken as a license to operate under, or a recommendation to infringe, any patents or patent applications.

© 2020 The Chemours Company FC, LLC. Glypure™ and any associated logos are trademarks or copyrights of The Chemours Company FC, LLC. Chemours™ and the Chemours Logo are trademarks of The Chemours Company.

Replaces: K-26051
C-10883 (3/20)